

Normal Families of Bicomplex Holomorphic Functions

K.S. Charak¹, D. Rochon², N. Sharma³

¹ Department of Mathematics, University of Jammu, Jammu-180 006, INDIA.
E-mail: kscharak7@rediffmail.com

² Département de mathématiques et d'informatique, Université du Québec à Trois-Rivières, C.P. 500 Trois-Rivières, Québec, Canada G9A 5H7.
E-mail: Dominic.Rochon@UQTR.CA, Web: www.3dfractals.com

³ Department of Mathematics, University of Jammu, Jammu-180 006, INDIA.
E-mail: narinder25sharma@sify.com

Abstract

In this article, we introduce the concept of normal families of bicomplex holomorphic functions to obtain a bicomplex Montel theorem. Moreover, we give a general definition of Fatou and Julia sets for bicomplex polynomials and we obtain a characterization of bicomplex Fatou and Julia sets in terms of Fatou set, Julia set and filled-in Julia set of one complex variable. Some 3D visual examples of bicomplex Julia sets are also given for the specific slice $\mathbf{j} = 0$.

Keywords: Bicomplex Numbers, Complex Clifford Algebras, Normal Families, Montel Theorem, Bicomplex Dynamics, Julia Set, Fatou Set.

1 Introduction

A family $\mathbf{F}$ of holomorphic functions defined on a domain $D \subseteq \mathbb{C}$ is said to be normal in D if every sequence in $\mathbf{F}$ has a subsequence converging uniformly on compact subsets of D to a function f . The limit function f is holomorphic on D (by Weierstrass Theorem) or the constant infinity. Various authors while studying the normality of a family of holomorphic functions take the limit function $f \neq \infty$ but for studying the normal families from complex dynamics point of view, one needs to include the case where the limit function $f \equiv \infty$. The former approach we shall call as restrictive approach while the later will be called the general approach towards normal families. The concept of normal families was introduced by P. Montel in 1907 [19]. For comprehensive account of normal families of meromorphic functions on domains in $\mathbb{C}$ one can refer to Joel Schiff's text [16], C.T. Chuang's text [20] and Zalcman's survey article [21]. With the renewed interest in normal families of meromorphic functions, arising largely from the important role they play in Complex Dynamics, it seems sensible to talk about normal families of holomorphic functions on different domains of different spaces thereby enabling one to study the dynamics of such functions. In this article we have considered the families of bicomplex holomorphic functions on bicomplex domains. Since this article lays the foundations of the subject **Normal Families of Bicomplex Holomorphic Functions** for future investigations in various possible directions, it is necessary to adopt a dual approach towards the study of normality of families of bicomplex holomorphic functions on bicomplex domains. The first approach is restrictive approach which gives rise to more interesting results when the normal families are studied in their own right. For example the converse of Montel Theorem holds under this approach. The second approach is the general approach in which though the converse of Montel Theorem fails to hold but is essentially required when the normal families are studied from the bicomplex dynamics point of view. During our discussions, we shall come across the situations where the differences lead to interesting conclusions. Besides complete discussion on Montel Theorem in various situations, we have defined Fatou, Julia and filled-in Julia sets of bicomplex polynomials and their representations in terms of their complex counterparts are obtained for the specific case of non-degenerate bicomplex polynomials of degree $d \geq 2$. Also, some 3D visual examples of bicomplex Julia sets are given for the specific slice $\mathbf{j} = 0$.

2 Preliminaries

2.1 Bicomplex Numbers

Bicomplex numbers are defined as

$$\mathbb{T} := \{z_1 + z_2 \mathbf{i}_2 \mid z_1, z_2 \in \mathbb{C}(\mathbf{i}_1)\} \quad (2.1)$$

where the imaginary units $\mathbf{i}_1, \mathbf{i}_2$ and $\mathbf{j}$ are governed by the rules: $\mathbf{i}_1^2 = \mathbf{i}_2^2 = -1$, $\mathbf{j}^2 = 1$ and

$$\begin{aligned}\mathbf{i}_1 \mathbf{i}_2 &= \mathbf{i}_2 \mathbf{i}_1 = \mathbf{j}, \\ \mathbf{i}_1 \mathbf{j} &= \mathbf{j} \mathbf{i}_1 = -\mathbf{i}_2, \\ \mathbf{i}_2 \mathbf{j} &= \mathbf{j} \mathbf{i}_2 = -\mathbf{i}_1.\end{aligned}\tag{2.2}$$

Note that we define $\mathbb{C}(\mathbf{i}_k) := \{x + y\mathbf{i}_k \mid \mathbf{i}_k^2 = -1 \text{ and } x, y \in \mathbb{R}\}$ for $k = 1, 2$. Hence, it is easy to see that the multiplication of two bicomplex numbers is commutative. In fact, the bicomplex numbers

$$\mathbb{T} \cong \text{Cl}_{\mathbb{C}}(1, 0) \cong \text{Cl}_{\mathbb{C}}(0, 1)$$

are **unique** among the *Complex Clifford algebras* (see [14]) in that they are commutative but not division algebra. It is also convenient to write the set of bicomplex numbers as

$$\mathbb{T} := \{w_0 + w_1 \mathbf{i}_1 + w_2 \mathbf{i}_2 + w_3 \mathbf{j} \mid w_0, w_1, w_2, w_3 \in \mathbb{R}\}.\tag{2.3}$$

In particular, in equation (2.1), if we put $z_1 = x$ and $z_2 = y\mathbf{i}_1$ with $x, y \in \mathbb{R}$, then we obtain the following subalgebra of hyperbolic numbers, also called duplex numbers (see, e.g. [12], [18]):

$$\mathbb{D} := \{x + y\mathbf{j} \mid \mathbf{j}^2 = 1, x, y \in \mathbb{R}\} \cong \text{Cl}_{\mathbb{R}}(0, 1).$$

Complex conjugation plays an important role both for algebraic and geometric properties of $\mathbb{C}$. For bicomplex numbers, there are three possible conjugations. Let $w \in \mathbb{T}$ and $z_1, z_2 \in \mathbb{C}(\mathbf{i}_1)$ such that $w = z_1 + z_2 \mathbf{i}_2$. Then we define the three conjugations as:

$$w^{\dagger_1} = (z_1 + z_2 \mathbf{i}_2)^{\dagger_1} := \bar{z}_1 + \bar{z}_2 \mathbf{i}_2,\tag{2.4a}$$

$$w^{\dagger_2} = (z_1 + z_2 \mathbf{i}_2)^{\dagger_2} := z_1 - z_2 \mathbf{i}_2,\tag{2.4b}$$

$$w^{\dagger_3} = (z_1 + z_2 \mathbf{i}_2)^{\dagger_3} := \bar{z}_1 - \bar{z}_2 \mathbf{i}_2,\tag{2.4c}$$

where $\bar{z}_k$ is the standard complex conjugate of complex number $z_k \in \mathbb{C}(\mathbf{i}_1)$. If we say that the bicomplex number $w = z_1 + z_2 \mathbf{i}_2 = w_0 + w_1 \mathbf{i}_1 + w_2 \mathbf{i}_2 + w_3 \mathbf{j}$ has the “signature” $(+++)$, then the conjugations of type 1, 2 or 3 of w have, respectively, the signatures $(+-+)$, $(++-)$ and $(+--+)$. We can verify easily that the composition of the conjugates gives the four-dimensional Klein group:

$\circ$	$\dagger_0$	$\dagger_1$	$\dagger_2$	$\dagger_3$
$\dagger_0$	$\dagger_0$	$\dagger_1$	$\dagger_2$	$\dagger_3$
$\dagger_1$	$\dagger_1$	$\dagger_0$	$\dagger_3$	$\dagger_2$
$\dagger_2$	$\dagger_2$	$\dagger_3$	$\dagger_0$	$\dagger_1$
$\dagger_3$	$\dagger_3$	$\dagger_2$	$\dagger_1$	$\dagger_0$

(2.5)

where $w^{\dagger_0} := w \ \forall w \in \mathbb{T}$.

The three kinds of conjugation all have some of the standard properties of conjugations, such as:

$$(s + t)^{\dagger_k} = s^{\dagger_k} + t^{\dagger_k}, \quad (2.6)$$

$$(s^{\dagger_k})^{\dagger_k} = s, \quad (2.7)$$

$$(s \cdot t)^{\dagger_k} = s^{\dagger_k} \cdot t^{\dagger_k}, \quad (2.8)$$

for $s, t \in \mathbb{T}$ and $k = 0, 1, 2, 3$.

We know that the product of a standard complex number with its conjugate gives the square of the Euclidean metric in $\mathbb{R}^2$. The analogs of this, for bicomplex numbers, are the following. Let $z_1, z_2 \in \mathbb{C}(\mathbf{i}_1)$ and $w = z_1 + z_2 \mathbf{i}_2 \in \mathbb{T}$, then we have that :

$$|w|_{\mathbf{i}_1}^2 := w \cdot w^{\dagger_2} = z_1^2 + z_2^2 \in \mathbb{C}(\mathbf{i}_1), \quad (2.9a)$$

$$|w|_{\mathbf{i}_2}^2 := w \cdot w^{\dagger_1} = (|z_1|^2 - |z_2|^2) + 2\operatorname{Re}(z_1 \bar{z}_2) \mathbf{i}_2 \in \mathbb{C}(\mathbf{i}_2), \quad (2.9b)$$

$$|w|_{\mathbf{j}}^2 := w \cdot w^{\dagger_3} = (|z_1|^2 + |z_2|^2) - 2\operatorname{Im}(z_1 \bar{z}_2) \mathbf{j} \in \mathbb{D}, \quad (2.9c)$$

where the subscript of the square modulus refers to the subalgebra $\mathbb{C}(\mathbf{i}_1), \mathbb{C}(\mathbf{i}_2)$ or $\mathbb{D}$ of $\mathbb{T}$ in which w is projected. Note that for $z_1, z_2 \in \mathbb{C}(\mathbf{i}_1)$ and $w = z_1 + z_2 \mathbf{i}_2 \in \mathbb{T}$, we can define the usual (Euclidean in $\mathbb{R}^4$) norm of w as $\|w\| = \sqrt{|z_1|^2 + |z_2|^2} = \sqrt{\operatorname{Re}(|w|_{\mathbf{j}}^2)}$.

It is easy to verify that $w \cdot \frac{w^{\dagger_2}}{|w|_{\mathbf{i}_1}^2} = 1$. Hence, the inverse of w is given by

$$w^{-1} = \frac{w^{\dagger_2}}{|w|_{\mathbf{i}_1}^2}. \quad (2.10)$$

From this, we find that the set $\mathcal{NC}$ of zero divisors of $\mathbb{T}$, called the *null-cone*, is given by $\{z_1 + z_2 \mathbf{i}_2 \mid z_1^2 + z_2^2 = 0\}$, which can be rewritten as

$$\mathcal{NC} = \{z(\mathbf{i}_1 \pm \mathbf{i}_2) \mid z \in \mathbb{C}(\mathbf{i}_1)\}. \quad (2.11)$$

2.2 Bicomplex Holomorphic Functions

It is also possible to define differentiability of a function at a point of $\mathbb{T}$:

Definition 1 *Let U be an open set of $\mathbb{T}$ and $w_0 \in U$. Then, $f : U \subseteq \mathbb{T} \longrightarrow \mathbb{T}$ is said to be $\mathbb{T}$ -differentiable at w_0 with derivative equal to $f'(w_0) \in \mathbb{T}$ if*

$$\lim_{\substack{w \rightarrow w_0 \\ (w - w_0 \text{ inv.})}} \frac{f(w) - f(w_0)}{w - w_0} = f'(w_0).$$

Note: *The subscript of the limit is there to recall that the division is possible only if $w - w_0$ is invertible.*

We also say that the function f is bicomplex holomorphic ($\mathbb{T}$ -holomorphic) on an open set U if and only if f is $\mathbb{T}$ -differentiable at each point of U . Using $w = z_1 + z_2 \mathbf{i}_2$, a bicomplex number w can be seen as an element (z_1, z_2) of $\mathbb{C}^2$, so a function $f(z_1 + z_2 \mathbf{i}_2) = f_1(z_1, z_2) + f_2(z_1, z_2) \mathbf{i}_2$ of $\mathbb{T}$ can be seen as a mapping $f(z_1, z_2) = (f_1(z_1, z_2), f_2(z_1, z_2))$ of $\mathbb{C}^2$. Here we have a characterization of such mappings:

Theorem 1 *Let U be an open set and $f : U \subseteq \mathbb{T} \longrightarrow \mathbb{T}$ such that $f \in C^1(U)$, and let $f(z_1 + z_2 \mathbf{i}_2) = f_1(z_1, z_2) + f_2(z_1, z_2) \mathbf{i}_2$. Then f is $\mathbb{T}$ -holomorphic on U if and only if*

$$f_1 \text{ and } f_2 \text{ are holomorphic in } z_1 \text{ and } z_2,$$

and

$$\frac{\partial f_1}{\partial z_1} = \frac{\partial f_2}{\partial z_2} \text{ and } \frac{\partial f_2}{\partial z_1} = -\frac{\partial f_1}{\partial z_2} \text{ on } U.$$

Moreover, $f' = \frac{\partial f_1}{\partial z_1} + \frac{\partial f_2}{\partial z_1} \mathbf{i}_2$ and $f'(w)$ is invertible if and only if $\det \mathcal{J}_f(w) \neq 0$.

This theorem can be obtained from results in [8] and [11]. Moreover, by the Hartogs theorem [17], it is possible to show that “ $f \in C^1(U)$ ” can be dropped from the hypotheses. Hence, it is natural to define the corresponding class of mappings for $\mathbb{C}^2$:

Definition 2 *The class of $\mathbb{T}$ -holomorphic mappings on a open set $U \subseteq \mathbb{C}^2$ is defined as follows:*

$$TH(U) := \{f : U \subseteq \mathbb{C}^2 \longrightarrow \mathbb{C}^2 \mid f \in H(U) \text{ and } \frac{\partial f_1}{\partial z_1} = \frac{\partial f_2}{\partial z_2}, \frac{\partial f_2}{\partial z_1} = -\frac{\partial f_1}{\partial z_2} \text{ on } U\}.$$

It is the subclass of holomorphic mappings of $\mathbb{C}^2$ satisfying the complexified Cauchy-Riemann equations.

We remark that $f \in TH(U)$ in terms of $\mathbb{C}^2$ if and only if f is $\mathbb{T}$ -differentiable on U . It is also important to know that every bicomplex number $z_1 + z_2 \mathbf{i}_2$ has the following unique idempotent representation:

$$z_1 + z_2 \mathbf{i}_2 = (z_1 - z_2 \mathbf{i}_1) \mathbf{e}_1 + (z_1 + z_2 \mathbf{i}_1) \mathbf{e}_2. \quad (2.12)$$

where $\mathbf{e}_1 = \frac{1+\mathbf{j}}{2}$ and $\mathbf{e}_2 = \frac{1-\mathbf{j}}{2}$. This representation is very useful because addition, multiplication and division can be done term-by-term. It is also easy to verify the following characterization of the non-invertible elements.

Proposition 1 *An element $w = z_1 + z_2 \mathbf{i}_2$ is in the null-cone if and only if $z_1 - z_2 \mathbf{i}_1 = 0$ or $z_1 + z_2 \mathbf{i}_1 = 0$.*

The notion of holomorphicity can also be seen with this kind of notation. For this we need to define the projections $\mathcal{P}_1, \mathcal{P}_2 : \mathbb{T} \longrightarrow \mathbb{C}(\mathbf{i}_1)$ as $\mathcal{P}_1(z_1 + z_2 \mathbf{i}_2) = z_1 - z_2 \mathbf{i}_1$ and $\mathcal{P}_2(z_1 + z_2 \mathbf{i}_2) = z_1 + z_2 \mathbf{i}_1$.

Definition 3 We say that $X \subseteq \mathbb{T}$ is a $\mathbb{T}$ -cartesian set determined by X_1 and X_2 if $X = X_1 \times_e X_2 := \{z_1 + z_2 \mathbf{i}_2 \in \mathbb{T} : z_1 + z_2 \mathbf{i}_2 = w_1 \mathbf{e}_1 + w_2 \mathbf{e}_2, (w_1, w_2) \in X_1 \times X_2\}$.

In [8] it is shown that if X_1 and X_2 are domains (open and connected) of $\mathbb{C}(\mathbf{i}_1)$ then $X_1 \times_e X_2$ is also a domain of $\mathbb{T}$. Then, a way to construct some “discus” (of center 0) in $\mathbb{T}$ is to take the $\mathbb{T}$ -cartesian product of two discs (of center 0) in $\mathbb{C}(\mathbf{i}_1)$. Hence, we define the “discus” with center $a = a_1 + a_2 \mathbf{i}_2$ of radius r_1 and r_2 of $\mathbb{T}$ as follows [8]: $D(a; r_1, r_2) := B^1(a_1 - a_2 \mathbf{i}_1, r_1) \times_e B^1(a_1 + a_2 \mathbf{i}_1, r_2) = \{z_1 + z_2 \mathbf{i}_2 : z_1 + z_2 \mathbf{i}_2 = w_1 \mathbf{e}_1 + w_2 \mathbf{e}_2, |w_1 - (a_1 - a_2 \mathbf{i}_1)| < r_1, |w_2 - (a_1 + a_2 \mathbf{i}_1)| < r_2\}$ where $B^1(z, r)$ is the open ball with center $z \in \mathbb{C}(\mathbf{i}_1)$ and radius $r > 0$. In the particular case where $r = r_1 = r_2$, $D(a; r, r)$ will be called the $\mathbb{T}$ -disc with center a and radius r . In particular, we define $\bar{D}(a; r_1, r_2) := \bar{B}^1(a_1 - a_2 \mathbf{i}_1, r_1) \times_e \bar{B}^1(a_1 + a_2 \mathbf{i}_1, r_2) \subset \bar{D}(a; r_1, r_2)$. We remark that $D(0; r, r)$ is, in fact, the Lie Ball (see [1]) of radius r in $\mathbb{T}$.

Now, it is possible to state the following striking theorems (see [8]):

Theorem 2 Let X_1 and X_2 be open sets in $\mathbb{C}(\mathbf{i}_1)$. If $f_{e1} : X_1 \rightarrow \mathbb{C}(\mathbf{i}_1)$ and $f_{e2} : X_2 \rightarrow \mathbb{C}(\mathbf{i}_1)$ are holomorphic functions of $\mathbb{C}(\mathbf{i}_1)$ on X_1 and X_2 respectively, then the function $f : X_1 \times_e X_2 \rightarrow \mathbb{T}$ defined as

$$f(z_1 + z_2 \mathbf{i}_2) = f_{e1}(z_1 - z_2 \mathbf{i}_1) \mathbf{e}_1 + f_{e2}(z_1 + z_2 \mathbf{i}_1) \mathbf{e}_2 \quad \forall z_1 + z_2 \mathbf{i}_2 \in X_1 \times_e X_2$$

is $\mathbb{T}$ -holomorphic on the open set $X_1 \times_e X_2$ and

$$f'(z_1 + z_2 \mathbf{i}_2) = f'_{e1}(z_1 - z_2 \mathbf{i}_1) \mathbf{e}_1 + f'_{e2}(z_1 + z_2 \mathbf{i}_1) \mathbf{e}_2$$

$$\forall z_1 + z_2 \mathbf{i}_2 \in X_1 \times_e X_2.$$

Theorem 3 Let X be an open set in $\mathbb{T}$, and let $f : X \rightarrow \mathbb{T}$ be a $\mathbb{T}$ -holomorphic function on X . Then there exist holomorphic functions $f_{e1} : X_1 \rightarrow \mathbb{C}(\mathbf{i}_1)$ and $f_{e2} : X_2 \rightarrow \mathbb{C}(\mathbf{i}_1)$ with $X_1 = \mathcal{P}_1(X)$ and $X_2 = \mathcal{P}_2(X)$, such that:

$$f(z_1 + z_2 \mathbf{i}_2) = f_{e1}(z_1 - z_2 \mathbf{i}_1) \mathbf{e}_1 + f_{e2}(z_1 + z_2 \mathbf{i}_1) \mathbf{e}_2$$

$$\forall z_1 + z_2 \mathbf{i}_2 \in X.$$

3 Bicomplex Montel Theorem

Since the concepts like **uniform boundedness**, **local uniform boundedness**, and **uniform convergence on compact sets** are defined for functions on any metric space and do not depend on bicomplex holomorphic functions, we assume these concepts in our discussion and refer the reader to any standard text on Analysis (e.g. see [4] and [13]). We start our discussion with the following definition of normality.

Definition 4 A family $\mathbf{F}$ of bicomplex holomorphic functions defined on a domain $D \subseteq \mathbb{T}$ is said to be **normal** in D if every sequence in $\mathbf{F}$ contains a subsequence which converges locally uniformly on D . $\mathbf{F}$ is said to be **normal at a point** $z \in D$ if it is normal in some neighborhood of z in D .

Let us consider $f : D \longrightarrow \mathbb{T}$ be a $\mathbb{T}$ -holomorphic function on D . Then by Theorem 3, there exist holomorphic functions $f_{e1} : \mathcal{P}_1(D) \longrightarrow \mathbb{C}(\mathbf{i}_1)$ and $f_{e2} : \mathcal{P}_2(D) \longrightarrow \mathbb{C}(\mathbf{i}_1)$ such that

$$f(z_1 + z_2 \mathbf{i}_2) = f_{e1}(z_1 - z_2 \mathbf{i}_1) \mathbf{e}_1 + f_{e2}(z_1 + z_2 \mathbf{i}_1) \mathbf{e}_2 \quad \forall z_1 + z_2 \mathbf{i}_2 \in D.$$

We define the **norm** of f on D as

$$\|f\| = \|f(z)\| = \left\{ \frac{|f_{e1}(z_1 - z_2 \mathbf{i}_1)|^2 + |f_{e2}(z_1 + z_2 \mathbf{i}_1)|^2}{2} \right\}^{\frac{1}{2}}, z = z_1 + z_2 \mathbf{i}_2 \in D.$$

One can easily see that

- $\|f\| \geq 0, \|f\| = 0$ iff $f \equiv 0$ on D ;
- $\|af\| = |a|\|f\|, a \in \mathbb{C}(\mathbf{i}_1)$;
- $\|f + g\| \leq \|f\| + \|g\|$;
- $\|fg\| \leq \sqrt{2}\|f\|\|g\|$.

Thus, the linear space of bicomplex holomorphic functions on a domain $D \subseteq \mathbb{T}$ is a normed space under the above norm.

We start with a **uniformly bounded** family $\mathbf{F}$ of bicomplex holomorphic functions. In this case, we can verify directly the following result.

Theorem 4 *A family $\mathbf{F}$ of bicomplex holomorphic functions defined on a bi-complex cartesian domain D is uniformly bounded on D if and only if $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is uniformly bounded on $\mathcal{P}_i(D)$, $i = 1, 2$.*

If we consider now a **locally uniformly bounded** family $\mathbf{F}$ of bicomplex holomorphic functions, we can prove a similar result since a set $K = \mathcal{P}_1(K) \times_e \mathcal{P}_2(K)$ is compact if and only if $\mathcal{P}_i(K)$ is compact for $i = 1, 2$.

Theorem 5 *A family $\mathbf{F}$ of bicomplex holomorphic functions defined on a bi-complex cartesian domain D is locally uniformly bounded on D if and only if $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is locally uniformly bounded on $\mathcal{P}_i(D)$, $i = 1, 2$.*

Proof Let $\mathbf{F}$ be locally uniformly bounded on D . Then for every compact set $K \subset D$ there is a constant $M(K)$ such that

$$\|f(z)\| \leq M, \quad \forall f \in \mathbf{F}, z = z_1 + z_2 \mathbf{i}_2 \in K.$$

Thus,

$$\left\{ \frac{|f_{e1}(z_1 - z_2 \mathbf{i}_1)|^2 + |f_{e2}(z_1 + z_2 \mathbf{i}_1)|^2}{2} \right\}^{\frac{1}{2}} \leq M, \quad \forall f_{ei} \in \mathbf{F}_{ei}, i = 1, 2$$

$$\forall z_1 - z_2 \mathbf{i}_1 \in \mathcal{P}_1(K), z_1 + z_2 \mathbf{i}_1 \in \mathcal{P}_2(K).$$

Therefore,

$$|f_{e1}(z_1 - z_2 \mathbf{i}_1)| \leq \sqrt{2}M, \forall f_{e1} \in \mathbf{F}_{e1}, \forall z_1 - z_2 \mathbf{i}_1 \in \mathcal{P}_1(K) \quad (3.1)$$

and

$$|f_{e2}(z_1 + z_2 \mathbf{i}_1)| \leq \sqrt{2}M, \forall f_{e2} \in \mathbf{F}_{e2} \forall z_1 + z_2 \mathbf{i}_1 \in \mathcal{P}_2(K). \quad (3.2)$$

Now, let K_1 be a compact subset of $\mathcal{P}_1(D)$. Then there is (always) a compact subset K_2 of $\mathcal{P}_2(D)$ (even singleton will do) such that $K_1 \times_e K_2 = K'$ say, is a compact subset of D with $\mathcal{P}_i(K') = K_i$, $i = 1, 2$. Thus (3.1) holds for any compact subset of $\mathcal{P}_1(D)$, and similarly for (3.2).

Conversely, suppose $\mathbf{F}_{ei}$ is locally uniformly bounded on $\mathcal{P}_i(D)$, $i = 1, 2$. Let K be any compact subset of D . Then by continuity of $\mathcal{P}_i$, $K_i = \mathcal{P}_i(K)$ is compact subset of $\mathcal{P}_i(D)$, $i = 1, 2$ and hence there are constants $M_1(K_1)$ and $M_2(K_2)$ such that

$$|f_{e1}(z_1 - z_2 \mathbf{i}_1)| \leq M_1, \forall f_{e1} \in \mathbf{F}_{e1}, \forall z_1 - z_2 \mathbf{i}_1 \in K_1$$

and

$$|f_{e2}(z_1 + z_2 \mathbf{i}_1)| \leq M_2, \forall f_{e2} \in \mathbf{F}_{e2} \forall z_1 + z_2 \mathbf{i}_1 \in K_2.$$

Therefore,

$$\|f(z)\| \leq \left\{ \frac{M_1^2 + M_2^2}{2} \right\}^{\frac{1}{2}}, \forall f \in \mathbf{F}, z = z_1 + z_2 \mathbf{i}_2 \in \mathcal{P}_1(K) \times_e \mathcal{P}_2(K). \quad (3.3)$$

Since $K \subseteq \mathcal{P}_1(K) \times_e \mathcal{P}_2(K)$, (3.3) holds for K also and this completes the proof. $\square$

What happens if D is not a bicomplex cartesian product? In the case of uniformly bounded family of bicomplex holomorphic functions (Theorem 4), it is easy to verify that the result is true for any domain. In the case of locally uniformly bounded family of bicomplex holomorphic functions, we need to recall the following results from the bicomplex function theory.

Remark 1 *A domain $D \subseteq \mathbb{T}$ is a domain of holomorphism for functions of a bicomplex variable if and only if D is a $\mathbb{T}$ -cartesian set ([8], Theorem 15.11), and if D is not a domain of holomorphism then $D \subsetneq \mathcal{P}_1(D) \times_e \mathcal{P}_2(D)$, and there exists a holomorphic function which is a bicomplex holomorphic continuation of the given function from D to $\mathcal{P}_1(D) \times_e \mathcal{P}_2(D)$ ([8] Corollary 15.4).*

Theorem 6 *A family $\mathbf{F}$ of bicomplex holomorphic functions defined on a arbitrary bicomplex domain D is locally uniformly bounded on D if and only if $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is locally uniformly bounded on $\mathcal{P}_i(D)$, $i = 1, 2$.*

Proof If $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is locally uniformly bounded on $\mathcal{P}_i(D)$ for $i = 1, 2$, from Remark 1, we can extend D to $\mathcal{P}_1(D) \times_e \mathcal{P}_2(D)$ and apply Theorem 5 to obtain that $\mathbf{F}$ is locally uniformly bounded on $\mathcal{P}_1(D) \times_e \mathcal{P}_2(D)$. For the other side, we need to recall that a family $\mathbf{F}$ is locally uniformly bounded on D if and only if the family $\mathbf{F}$ is **locally bounded** on D i.e. for each $w_0 \in D$ there is a positive

number $M = M(w_0)$ and a neighbourhood $D(w_0; r, r) \subset D$ such that $\|f(w)\| \leq M$ for all $w \in D(w_0; r, r)$ and all $f \in \mathbf{F}$ (see [16]). Since $D(w_0; r, r) \subset D$ is a bicomplex cartesian product of two discs in the plane, it is easy to verify that the family $\mathbf{F}_{ei}$ is bounded by $\sqrt{2}M(w_0)$ on $D(\mathcal{P}_i(w_0), r) \subset \mathcal{P}_i(D)$ for $i = 1, 2$. As w_0 was arbitrary, $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is locally bounded on $\mathcal{P}_i(D)$, $i = 1, 2$. $\square$

We are now ready to prove the bicomplex version of the Montel theorem.

Lemma 1 *Let $\mathbf{F}$ be a family of bicomplex holomorphic functions defined on a bicomplex domain D . If $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is normal on $\mathcal{P}_i(D)$ for $i = 1, 2$ then $\mathbf{F}$ is normal on D .*

Proof Suppose that $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is normal on $\mathcal{P}_i(D) = D_i$, $i = 1, 2$. We want to show that $\mathbf{F}$ is normal in D . Let $\{F_n\}$ be any sequence in $\mathbf{F}$ and K be any compact subset of D . Then $\{\mathcal{P}_1(F_n)\} = \{(f_n)_1\}$ is a sequence in $\mathbf{F}_{e1} = \mathcal{P}_1(\mathbf{F})$. Since $\mathbf{F}_{e1} = \mathcal{P}_1(\mathbf{F})$ is normal in $\mathcal{P}_1(D)$ then $\{(f_n)_1\}$ has a subsequence $\{(f_{n_k})_1\}$ which converges uniformly on $\mathcal{P}_1(K)$ to a $\mathbb{C}(\mathbf{i}_1)$ -function. Now, consider $\{F_{n_k}\}$ in $\mathbf{F}$. Then $\{\mathcal{P}_2(F_{n_k})\} = \{(f_{n_k})_2\}$ is a sequence in $\mathbf{F}_{e2} = \mathcal{P}_2(\mathbf{F})$. Since $\mathbf{F}_{e2} = \mathcal{P}_2(\mathbf{F})$ is normal in $\mathcal{P}_2(D)$ then $\{(f_{n_k})_2\}$ has a subsequence $\{(f_{n_{k_l}})_2\}$ which converges uniformly on $\mathcal{P}_2(K)$ to a $\mathbb{C}(\mathbf{i}_1)$ -function. This implies that $\{(f_{n_{k_l}})_1 \mathbf{e}_1 + (f_{n_{k_l}})_2 \mathbf{e}_2\}$ is a subsequence of $\{F_n\}$ which converges uniformly on $\mathcal{P}_1(K) \times_e \mathcal{P}_2(K) \supseteq K$ to a bicomplex function showing that $\mathbf{F}$ is normal in D . $\square$

Theorem 7 (Montel) *Every locally uniformly bounded family of bicomplex holomorphic functions defined on a bicomplex domain is a normal family.*

Proof Let $\mathbf{F}$ be a locally uniformly bounded family of bicomplex holomorphic functions defined on a domain $D \subseteq \mathbb{T}$. Using Theorem 6, we have that $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is locally uniformly bounded on $\mathcal{P}_i(D)$, $i = 1, 2$. Hence, from the classical Montel Theorem, $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is normal on $\mathcal{P}_i(D)$ for $i = 1, 2$ and by Lemma 1 we obtain that $\mathbf{F}$ is normal on D . $\square$

Note: The converse of Bicomplex Montel Theorem is also true. Indeed, suppose that $\mathbf{F}$ is normal and not locally uniformly bounded in D . Then in some closed disc $\overline{D}(a; r_1, r_2)$ in the domain D , for each $n \in \mathbb{N}$ there is a function $f_n \in \mathbf{F}$ and a point $w_n \in \overline{D}(a; r_1, r_2)$ such that $\|f_n(w_n)\| > n$. Since $\mathbf{F}$ is normal, there is a subsequence $\{f_{n_k}\}$ of $\{f_n\}$ converging uniformly on $\overline{D}(a; r_1, r_2)$ to a bicomplex (holomorphic) function f . That is, for some positive integer n_0 , we have

$$\|f_{n_k}(w) - f(w)\| < 1, \quad \forall k \geq n_0, \text{ and } w \in \overline{D}(a; r_1, r_2).$$

Thus, if $M = \max_{z \in \overline{D}(a; r_1, r_2)} \|f(w)\|$, then $\|f_{n_k}(w)\| \leq 1 + M$, $\forall w \in \overline{D}(a; r_1, r_2)$ and this is a contradiction.

The above discussion permits to establish the following results.

Theorem 8 *The family $\mathbf{F}$ of bicomplex holomorphic functions is normal on the arbitrary domain D if and only if $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is normal on $\mathcal{P}_i(D)$ for $i = 1, 2$.*

Corollary 1 *If the family $\mathbf{F}$ of bicomplex holomorphic functions is normal on a arbitrary domain $D \neq \mathcal{P}_1(D) \times_e \mathcal{P}_2(D)$, then $\mathbf{F}$ is normal on the larger domain $\mathcal{P}_1(D) \times_e \mathcal{P}_2(D)$.*

Corollary 2 *A family of bicomplex holomorphic functions $\mathbf{F}$ is normal on a arbitrary domain D if and only if $\mathbf{F}$ is normal at each point of D .*

4 Bicomplex Montel Theorem from Montel Theorem of $\mathbb{C}^2$

In this section, we want to show that it is possible to see the Bicomplex Montel Theorem (Theorem 7) as a particular case of the following Montel theorem of several complex variables (see [15]).

Theorem 9 *Let $D \subset \mathbb{C}^n$ be an open set and $\mathbf{F} \subset \mathcal{O}(D, \mathbb{C}^n)$ be a family of holomorphic mappings. Then the following are equivalent:*

1. *The family $\mathbf{F}$ is locally uniformly bounded.*
2. *The family $\mathbf{F}$ is relatively compact in $\mathcal{O}(D, \mathbb{C}^n)$.*

Since, $TH(D) \subset \mathcal{O}(D, \mathbb{C}^2)$, we obtain directly the desired result using the fact that a family $\mathbf{F}$ is relatively compact in $\mathcal{O}(D, \mathbb{C}^n)$, i.e. $\overline{\mathbf{F}}$ is compact in $\mathcal{O}(D, \mathbb{C}^n)$, if and only if $\mathbf{F}$ is a normal family (see [4]). Moreover, Theorem 9 will be proven for the specific class $TH(D)$ instead of $\mathcal{O}(D, \mathbb{C}^2)$ if we can show that $TH(D)$ is closed in $\mathcal{O}(D, \mathbb{C}^2)$ with the compact convergence topology. This is a direct consequence of the following **Bicomplex Weierstrass Theorem**.

Lemma 2 *Let $\{f_n\}$ be a sequence of bicomplex holomorphic functions which converges locally uniformly to a function f on a $\mathbb{T}$ -disc $D(a_1 + a_2 \mathbf{i}_2; r, r)$. Then f is bicomplex holomorphic in $D(a_1 + a_2 \mathbf{i}_2; r, r)$.*

Proof Since $f_n(z_1 + z_2 \mathbf{i}_2)$ is $\mathbb{T}$ -holomorphic on $D(a_1 + a_2 \mathbf{i}_2; r, r) \forall n \in \mathbb{N}$, we have from Theorem 3 that

$$(f_{ei})_n : \mathcal{P}_i(D(a_1 + a_2 \mathbf{i}_2, r)) \longrightarrow \mathbb{C}(\mathbf{i}_1)$$

is holomorphic for $i = 1, 2, \forall n \in \mathbb{N}$. Since $D(a_1 + a_2 \mathbf{i}_2; r, r)$ is a bicomplex cartesian product, by the Weierstrass theorem of one complex variable, the sequence $(f_{ei})_n$ must converges locally uniformly to the holomorphic function f_{ei} on $D(\mathcal{P}_i(a_1 + a_2 \mathbf{i}_2), r)$ for $i = 1, 2$. Therefore, from Theorem 2, the function $f(z_1 + z_2 \mathbf{i}_2) = f_{e1}(z_1 - z_2 \mathbf{i}_1) \mathbf{e}_1 + f_{e2}(z_1 + z_2 \mathbf{i}_1) \mathbf{e}_2$ is $\mathbb{T}$ -holomorphic on $D(a_1 + a_2 \mathbf{i}_2; r, r)$. $\square$

Theorem 10 (Weierstrass) *Let $\{f_n\}$ be a sequence of bicomplex holomorphic functions on a domain D which converges uniformly on compact subsets of D to a function f . Then f is bicomplex holomorphic in D .*

Proof For an arbitrary $w_0 \in D$, choose a $\mathbb{T}$ -disc $D(w_0; r, r) \subset D$. Since $f_n(w) \rightarrow f(w)$ locally uniformly on D , by Lemma 2, f is $\mathbb{T}$ -holomorphic on D . As w_0 was arbitrary, $f(w)$ is $\mathbb{T}$ -holomorphic on D . $\square$

5 A More General Definition of Normality

To carry further the study of normal families of bicomplex holomorphic functions particularly to consider the dynamics of bicomplex holomorphic functions, we propose the following more general definition of normality.

Definition 5 *A family $\mathbf{F}$ of bicomplex holomorphic functions defined on a domain $D \subseteq \mathbb{T}$ is said to be **normal** in D if every sequence in $\mathbf{F}$ contains a subsequence which on compact subsets of D either converges uniformly to a limit function or converges uniformly to ∞ . $\mathbf{F}$ is said to be **normal at a point** $z \in D$ if it is normal in some neighborhood of z in D .*

Remark 2 *We say that a sequence $\{w_n\}$ of bicomplex numbers converges to ∞ if and only if the norm $\{\|w_n\|\}$ converges to ∞ .*

We note that our proofs of the Bicomplex Montel Theorem work also in this situation. However, as for one complex variable, the converse of Theorem 7 will not remain valid with this more complete definition of normality (see [16]).

Remark 3 *Both the situations in the last definition may occur simultaneously. For example, consider the family $\{R^n(w) \mid R(w) = w^2 \text{ and } n \in \mathbb{N}\}$ of bicomplex holomorphic functions on $\mathbb{T}$. Then, by using the idempotent representation and results from one complex variable theory of normal families, we find that this family is normal on $A \cup B$, where*

$$A = \{w = w_1 \mathbf{e}_1 + w_2 \mathbf{e}_2 : |w_1| < 1, |w_2| < 1\}$$

and

$$B = \{w = w_1 \mathbf{e}_1 + w_2 \mathbf{e}_2 : |w_1| > 1, |w_2| > 1\}$$

On the set A , normality is under the first situation whereas on the set B the normality is under the second situation.

Example 1 *Consider the family*

$$\mathbf{F} = \{f_n(w) = nw : w = z_1 + z_2 \mathbf{i}_2, \quad n \in \mathbb{Z}\}.$$

Then $f_n(0) \rightarrow 0$, but $f_n(w) \rightarrow \infty$ for $w \neq 0$. It follows that $\mathbf{F}$ cannot be normal in any domain containing the origin.

Now, let us prove that Theorem 8 is only true in one direction with this more general definition of normality.

Theorem 11 *Let $\mathbf{F}$ be a family of bicomplex holomorphic functions defined on a domain D . If $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is normal on $\mathcal{P}_i(D)$ for $i = 1, 2$ then $\mathbf{F}$ is normal on D .*

Proof Suppose that $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is normal on $\mathcal{P}_i(D) = D_i$, $i = 1, 2$. We want to show that $\mathbf{F}$ is normal in D . Let $\{F_n\}$ be any sequence in $\mathbf{F}$ and K be any compact subset of D . Then $\{\mathcal{P}_1(F_n)\} = \{(f_n)_1\}$ is a sequence in $\mathbf{F}_{e1} = \mathcal{P}_1(\mathbf{F})$. Since $\mathbf{F}_{e1} = \mathcal{P}_1(\mathbf{F})$ is normal in $\mathcal{P}_1(D)$ then $\{(f_n)_1\}$ has a subsequence $\{(f_{n_k})_1\}$ which converges uniformly on $\mathcal{P}_1(K)$ to either a $\mathbb{C}(\mathbf{i}_1)$ -function or to ∞ . Now, consider $\{F_{n_k}\}$ in $\mathbf{F}$. Then $\{\mathcal{P}_2(F_{n_k})\} = \{(f_{n_k})_2\}$ is a sequence in $\mathbf{F}_{e2} = \mathcal{P}_2(\mathbf{F})$. Since $\mathbf{F}_{e2} = \mathcal{P}_2(\mathbf{F})$ is normal in $\mathcal{P}_2(D)$ then $\{(f_{n_k})_2\}$ has a subsequence $\{(f_{n_{k_l}})_2\}$ which converges uniformly on $\mathcal{P}_2(K)$ to either a $\mathbb{C}(\mathbf{i}_1)$ -function or to ∞ . This implies that $\{(f_{n_{k_l}})_1 \mathbf{e}_1 + (f_{n_{k_l}})_2 \mathbf{e}_2\}$ is a subsequence of $\{F_n\}$ which converges uniformly on $\mathcal{P}_1(K) \times_e \mathcal{P}_2(K) \supseteq K$ to a bicomplex function or to ∞ showing that $\mathbf{F}$ is normal in D . $\square$

Here is the counterexample for the other side.

Example 2 Let X_1 and X_2 be domains in $\mathbb{C}(\mathbf{i}_1)$ containing the origin. Let $D = (X_1 \times_e X_2) - \{0\}$. Then D is not a bicomplex cartesian domain because $D \neq \mathcal{P}_1(D) \times_e \mathcal{P}_2(D)$. Now the family

$$\mathbf{F} = \{nw : w = z_1 + z_2 \mathbf{i}_2, \quad n \in \mathbb{N}\}$$

is normal in the domain D (by the proposed definition of normality as above) but $\mathbf{F}_{ei} = \mathcal{P}_i(\mathbf{F})$ is not normal in $\mathcal{P}_i(D)$, $i = 1, 2$ as it contains the origin.

Moreover, the next examples show that the converse of Theorem 11 is not true even if the domain D is a bicomplex cartesian product.

Example 3 Consider the family

$$\mathcal{F} = \{nz : z \in \mathbb{C}(\mathbf{i}_1), \quad n \in \mathbb{Z}\}$$

on $\mathbb{C}(\mathbf{i}_1)$. Then $\mathcal{F}$ is normal on the punctured disc $D(0, 1) - \{0\} \subset \mathbb{C}(\mathbf{i}_1)$ but not normal on the disc $D(0, 1) \subset \mathbb{C}(\mathbf{i}_1)$. However, the bicomplex family

$$\mathbf{F} := \mathbf{F}_{e1} \mathbf{e}_1 + \mathbf{F}_{e2} \mathbf{e}_2 = \{nw : w = z_1 + z_2 \mathbf{i}_2, \quad n \in \mathbb{N}\}$$

where $\mathbf{F}_{e1} = \mathcal{F}$ is normal in the following bicomplex cartesian product:

$$(D(0, 1) - \{0\}) \times_e D(0, 1)$$

since the limit function is identically infinite.

Example 4 Consider the family

$$\mathcal{F} = \{R^n(z) \mid R(z) = z^2 \text{ and } n \in \mathbb{N}\}$$

on $\mathbb{C}(\mathbf{i}_1)$. Then $\mathcal{F}$ is normal on $D_1 = \{z : |z| > 1\} \subset \mathbb{C}(\mathbf{i}_1)$ where here the limit function is identically infinite, but not normal on $\mathbb{C}(\mathbf{i}_1)$ since $\{|z| = 1\} \subset \mathbb{C}(\mathbf{i}_1)$. However, the bicomplex family

$$\mathbf{F} := \mathbf{F}_{e1} \mathbf{e}_1 + \mathbf{F}_{e2} \mathbf{e}_2 = \{R^n(w) \mid R(w) = w^2 \text{ and } n \in \mathbb{N}\}$$

where $\mathbf{F}_{e1} = \mathbf{F}_{e2} = \mathcal{F}$, is normal in the following bicomplex cartesian product:

$$D_1 \times_e \mathbb{C}(\mathbf{i}_1)$$

since the limit function is identically infinite.

6 Foundation of Bicomplex Dynamics: Fatou and Julia Sets for Polynomials

We conclude this article with the following general definition of **Fatou** and **Julia** sets for bicomplex polynomials.

Definition 6 *Let $P(\zeta)$ be a bicomplex polynomials. We define the bicomplex Julia set for P as*

$$\mathcal{J}_2(P) = \{\zeta \in \mathbb{T} \mid \{P^{on}(\zeta)\} \text{ is not normal}\} \quad (6.1)$$

and the bicomplex Fatou (or stable) set as

$$\mathcal{F}_2(P) = \mathbb{T} - \mathcal{J}_2(P). \quad (6.2)$$

Hence, about each point $\zeta \in \mathcal{F}_2(P)$, there is a neighborhood N_ζ in which $\{P^{on}(\zeta)\}$ is a normal family. Therefore, $\mathcal{F}_2(P)$ is an open set, the connected components of which are the maximal domains of normality of $\{P^{on}(\zeta)\}$, and $\mathcal{J}_2(P)$ is a closed set.

From Theorem 11, we obtain the following inclusion:

$$\mathcal{J}_2(P) \subset \{z_1 + z_2 \mathbf{i}_2 \in \mathbb{T} \mid \{[\mathcal{P}_1(P)]^{on}(z_1 - z_2 \mathbf{i}_1)\} \text{ or} \quad (6.3)$$

$$\{[\mathcal{P}_2(P)]^{on}(z_1 + z_2 \mathbf{i}_1)\} \text{ is not normal} \} \quad (6.4)$$

$$= [\mathcal{J}_1(\mathcal{P}_1(P)) \times_e \mathbb{C}(\mathbf{i}_1)] \cup [\mathbb{C}(\mathbf{i}_1) \times_e \mathcal{J}_1(\mathcal{P}_2(P))]. \quad (6.5)$$

However, from Example 4, we know that (6.3) cannot be transformed into equality. In fact, to obtain a characterization of bicomplex Julia sets in terms of one variable dynamics we need to use the concept of filled-in Julia set. As for the complex case, the bicomplex **filled-in Julia** set $K_2(P)$ of a polynomial P is defined as the set of all points ζ of dynamical space that have bounded orbit with respect to P , that is to say:

$$\mathcal{K}_2(P) = \{\zeta \in \mathbb{T} \mid \{P^{on}(\zeta)\} \nrightarrow \infty \text{ as } n \rightarrow \infty\}. \quad (6.6)$$

We remark that $\mathcal{K}_2(P)$ is a closed set.

As for the classical case (see [3], P.65), we need to consider polynomials of degree $d \geq 2$ to be able to see a bicomplex Julia set as the boundary of a bicomplex filled-in Julia set. In fact, to decompose $P(w)$ in terms of two complex polynomials of degree $d \geq 2$, we must also consider non-degenerate bicomplex polynomials of the form $P(w) = a_d w^d + a_{d-1} w^{d-1} + \dots + a_0$ where $a_d \notin \mathcal{NC}$.

Under these specifications, we have the following result.

Theorem 12 *Let $P(\zeta)$ be a non-degenerate bicomplex polynomials of degree $d \geq 2$. Then,*

$$\partial \mathcal{K}_2(P) = \mathcal{J}_2(P). \quad (6.7)$$

Now, using the concept of normality in terms of **Definition 4**, we obtain the following characterization of $\mathcal{K}_2(P) - \mathcal{J}_2(P)$:

$$\mathcal{K}_2(P) - \mathcal{J}_2(P) = \{\zeta \in \mathbb{T} \mid \{P^{on}(\zeta)\} \text{ is normal}\}. \quad (6.8)$$

Moreover, using the idempotent representation, it is easy to see that the bicomplex filled-in Julia set $\mathcal{K}_2(P)$ can be expressed in terms of two filled-in Julia sets in the plane. More specifically,

$$\mathcal{K}_2(P) = \mathcal{K}_1(\mathcal{P}_1(P)) \times_e \mathcal{K}_1(\mathcal{P}_2(P)). \quad (6.9)$$

Hence, since $\partial[\mathcal{K}_1(\mathcal{P}_1(P)) \times_e \mathcal{K}_1(\mathcal{P}_2(P))] = [\partial \mathcal{K}_1(\mathcal{P}_1(P)) \times_e \mathcal{K}_1(\mathcal{P}_2(P))] \cup [\mathcal{K}_1(\mathcal{P}_1(P)) \times_e \partial \mathcal{K}_1(\mathcal{P}_2(P))]$, we have the following characterization of the bicomplex Julia set $\mathcal{J}_2(P)$ in terms of one complex variable dynamics.

Theorem 13 *Let $P(\zeta)$ be a non-degenerate bicomplex polynomials of degree $d \geq 2$. Then,*

$$\mathcal{J}_2(P) = [\mathcal{J}_1(\mathcal{P}_1(P)) \times_e \mathcal{K}_1(\mathcal{P}_2(P))] \cup [\mathcal{K}_1(\mathcal{P}_1(P)) \times_e \mathcal{J}_1(\mathcal{P}_2(P))]. \quad (6.10)$$

Example 5 *Consider the bicomplex polynomial:*

$$P(w) = w^2.$$

We can verify that $\mathcal{P}_k(w^2) = z^2$ for $k = 1, 2$. In the complex plane (in $\mathbf{i}_1$), it is well known that $\mathcal{K}_1(z^2) = \{z : |z| < 1\}$ and $\mathcal{J}_1(z^2) = \{z : |z| = 1\}$ (see [2] or [16]). Hence, using Theorem 13, we obtain that

$$\mathcal{J}_2(P(w)) = [\{z : |z| = 1\} \times_e \{z : |z| < 1\}] \cup [\{z : |z| < 1\} \times_e \{z : |z| = 1\}]. \quad (6.11)$$




Figure 1: $c = (0.27)\mathbf{e}_1 + (0.27)\mathbf{e}_2$

Remark 4 In the particular case of the bicomplex quadratic polynomial

$$P_c(\zeta) = \zeta^2 + c, \quad (6.12)$$

the **fundamental** definition of bicomplex Julia set of this article (see 6.1) coincide with the definition, using boundary of bicomplex filled-in Julia set, introduced by D. Rochon in [9, 10] (see Theorem 12). Moreover, using some distance estimation formulas that can be used to ray traced slices of bicomplex filled-in Julia sets in dimension three (see [7]), we obtain some visual examples (see Fig. 1, 2, 3 and 4) of bicomplex Julia sets $\mathcal{K}_2(P_c)$ for the specific slice $\mathbf{j} = 0$.

Example 6 Consider the bicomplex polynomial:

$$P(w) = w^2 + 0.27.$$

We can verify that $\mathcal{P}_k(w^2) = z^2 + 0.27$ for $k = 1, 2$. In the complex plane (in $\mathbf{i}_1$), it is well known that $\mathcal{K}_1(z^2 + 0.27) = \mathcal{J}_1(z^2 + 0.27)$ is a Cantor set. We shall denote such Cantor set by $\mathcal{C}_{0.27}$. Hence, using Theorem 13, we obtain that

$$\mathcal{J}_2(P) = \mathcal{C}_{0.27} \times_e \mathcal{C}_{0.27} \quad (\text{Fig. 1}). \quad (6.13)$$

Example 7 Consider the bicomplex polynomial:

$$P(w) = w^2 - 1.754878.$$

We can verify that $\mathcal{P}_k(w^2) = z^2 - 1.754878$ for $k = 1, 2$. In the complex plane (in $\mathbf{i}_1$), it is well known that $\mathcal{K}_1(z^2 - 1.754878)$ is the so-called **Airplane**. We shall denote this set by $\mathcal{A}$. Hence, using Theorem 13, we obtain that

$$\mathcal{J}_2(P) = [\partial\mathcal{A} \times_e \mathcal{A}] \cup [\mathcal{A} \times_e \partial\mathcal{A}] \quad (\text{Fig. 2}). \quad (6.14)$$

Example 8 Consider the bicomplex polynomial:

$$P(w) = w^2 + [(0.26)\mathbf{e}_1 + (-1.754878)\mathbf{e}_2].$$

We can verify that $\mathcal{P}_1(w^2 + [(0.26)\mathbf{e}_1 + (-1.754878)\mathbf{e}_2]) = z^2 + 0.26$ and $\mathcal{P}_1(w^2 + [(0.26)\mathbf{e}_1 + (-1.754878)\mathbf{e}_2]) = z^2 - 1.754878$. Hence, using Theorem 13, we obtain that

$$\mathcal{J}_2(P) = \mathcal{C}_{0.26} \times_e \mathcal{A} \quad (\text{Fig. 3}). \quad (6.15)$$




Figure 2: $c = (-1.754878)\mathbf{e}_1 + (-1.754878)\mathbf{e}_2$



Figure 3: $c = (0.26)\mathbf{e}_1 + (-1.754878)\mathbf{e}_2$

Example 9 Consider the bicomplex polynomial:

$$P(w) = w^2 + [(-0.123 + 0.745\mathbf{i}_1)\mathbf{e}_1 + (-0.391 - 0.587\mathbf{i}_1)\mathbf{e}_2].$$

We can verify that $\mathcal{P}_1(w^2 + [(-0.123 + 0.745\mathbf{i}_1)\mathbf{e}_1 + (-0.391 - 0.587\mathbf{i}_1)\mathbf{e}_2]) = z^2 + (-0.123 + 0.745\mathbf{i}_1)$ and $\mathcal{P}_1(w^2 + [(-0.123 + 0.745\mathbf{i}_1)\mathbf{e}_1 + (-0.391 - 0.587\mathbf{i}_1)\mathbf{e}_2]) = z^2 + (-0.391 - 0.587\mathbf{i}_1)$. In the complex plane (in $\mathbf{i}_1$), it is well known that $\mathcal{D} := \mathcal{K}_1(z^2 + (-0.123 + 0.745\mathbf{i}_1))$ is the so-called **Douady's Rabbit** and $\mathcal{S} := \mathcal{K}_1(z^2 + (-0.391 - 0.587\mathbf{i}_1))$ is a **Siegel Disk**. Hence, using Theorem 13, we obtain that

$$\mathcal{J}_2(P) = [\partial\mathcal{D} \times_e \mathcal{S}] \cup [\mathcal{D} \times_e \partial\mathcal{S}] \quad (\text{Fig. 4}). \quad (6.16)$$

Remark 5 By using the definition of bicomplex Fatou set as the complement of bicomplex Julia set (6.10) leads us to characterize the bicomplex Fatou set of non-degenerate bicomplex polynomials of degree $d \geq 2$ as

$$\begin{aligned} \mathcal{F}_2(P) = & [\mathcal{F}_1(\mathcal{P}_1(P)) \times_e \mathcal{F}_1(\mathcal{P}_2(P))] \cup [\mathcal{F}_1(\mathcal{P}_1(P))_\infty \times_e \mathcal{J}_1(\mathcal{P}_2(P))] \\ & \cup [\mathcal{J}_1(\mathcal{P}_1(P)) \times_e \mathcal{F}_1(\mathcal{P}_2(P))_\infty] \end{aligned} \quad (6.17)$$

where $\mathcal{F}_1(\mathcal{P}_i(P))_\infty$, $i = 1, 2$ denotes the unbounded component of the Fatou set of projections of P .




Figure 4: $c = (-0.123 + 0.745\mathbf{i}_1)\mathbf{e}_1 + (-0.391 - 0.587\mathbf{i}_1)\mathbf{e}_2$

Example 10 Consider the bicomplex polynomial:

$$P(w) = w^2.$$

In the complex plane (in $\mathbf{i}_1$), it is well known that

$$\mathcal{F}_1(z^2) = \{z : |z| < 1\} \cup \{z : |z| > 1\}$$

and $\mathcal{F}_1(z^2)_\infty = \{z : |z| > 1\}$. Hence, using (6.17), we obtain that

$$\begin{aligned} \mathcal{F}_2(P) = & [(\{z : |z| < 1\} \cup \{z : |z| > 1\}) \times_e (\{z : |z| < 1\} \cup \{z : |z| > 1\})] \\ & \cup [\{z : |z| > 1\} \times_e \{z : |z| = 1\}] \\ & \cup [\{z : |z| = 1\} \times_e \{z : |z| > 1\}]. \end{aligned} \quad (6.18)$$

As a direct consequence of Theorem 13, the bicomplex Julia set $\mathcal{J}_2(P)$ is completely invariant under the substitution $(w, P(w))$ when P is a non-degenerate bicomplex polynomials of degree $d \geq 2$. The next theorem will prove this result in general.

Definition 7 Let $f(z_1 + z_2\mathbf{i}_2) = f_{e1}(z_1 - z_2\mathbf{i}_1)\mathbf{e}_1 + f_{e1}(z_1 + z_2\mathbf{i}_1)\mathbf{e}_2 : D \longrightarrow \mathbb{T}$ be any bicomplex function. The function f is said to be strongly non-constant on D if f_{ei} is non-constant on $\mathcal{P}_i(D)$ for $i = 1, 2$.

Theorem 14 Let f be an entire strongly non-constant $\mathbb{T}$ -holomorphic function and

$$\mathcal{J}_2(f) := \{\zeta \in \mathbb{T} \mid \{f^{\text{on}}(\zeta)\} \text{ is not normal}\}.$$

Then,

1. If a point $w_0 \in \mathcal{J}_2(f)$, then $f(w_0) \in \mathcal{J}_2(f)$;
2. If $w_0 \in \mathcal{J}_2(f)$ and w_1 is a point such that $f(w_1) = w_0$ then $w_1 \in \mathcal{J}_2(f)$.

Proof Since our definition of normality (Def. 5) is analogous to the related notion in the complex plane, the proof of the theorem is same as the proof of the corresponding result in the plane (see [5], Theorem 2.17 and 2.18). **Note:** The condition for f to be strongly non-constant is needed in the proof of (2.) to be able to use the open mapping theorem in each components. $\square$

Acknowledgements

The research of D.R. is partly supported by grants from CRSNG of Canada and FQRNT of Québec.

References

- [1] V. Avaniessian, *Cellule d'harmonicit  et prolongement analytique complexe*, (Travaux en cours, Paris, 1985).
- [2] A. F. Beardon, *Iteration of Rational Functions*, Springer-Verlag, New York, (1991).
- [3] L. Carleson and T.W. Gamelin, *Complex Dynamics*, Springer-Verlag, New York (1993).
- [4] J. B. Conway, *Functions of one complex variables I*, Springer-Verlag, New York (1978).
- [5] C.T. Chuang and C.C. Yang, *Fix-points and Factorization of meromorphic functions*, World Scientific (1990).
- [6] N. Fleury, M. Rausch de Traubenberg and R.M. Yamaleev, Commutative extended complex numbers and connected trigonometry, *J. Math. Ann. and Appl.* **180**, 431–457 (1993).
- [7]  . Martineau and D. Rochon, On a Bicomplex Distance Estimation for the Tetraebrot, *International Journal of Bifurcation and Chaos*, **15**(9), 1–12 (2005).
- [8] G.B. Price, *An introduction to multicomplex spaces and functions*, Marcel Dekker Inc., New York (1991).
- [9] D. Rochon, A generalized Mandelbrot set for bicomplex numbers, *Fractal* **8**, 355–368 (2000).
- [10] D. Rochon, On a Generalized Fatou-Julia Theorem, *Fractals* **11**(3), 213–219 (2003).
- [11] D. Rochon, Sur une g n ralisation des nombres complexes: les t tranombres, (M. Sc. Universit  de Montr al, 1997).

- [12] D. Rochon and M. Shapiro, On algebraic properties of bicomplex and hyperbolic numbers, *Anal. Univ. Oradea*, fasc. math., **11**, 71–110 (2004).
- [13] W. Rudin, *Real and Complex Analysis 3rd ed.*, McGraw-Hill, New York (1976).
- [14] J. Ryan, Complexified Clifford Analysis, *Complex Variables* **1**, 119–149 (1982).
- [15] V. Scheidemann, *Introduction to Complex Analysis in Several Variables*, Birkhäuser Verlag, Basel (2005).
- [16] J.L. Schiff, *Normal Families*, Springer-Verlag, New York (1993).
- [17] B.V. Shabat, *Introduction to Complex Analysis part II: Functions of Several Variables*, American Mathematical Society (1992).
- [18] G. Sobczyk, The hyperbolic number plane, *Coll. Maths. Jour.* **26**, No. 4, 268–280 (1995).
- [19] P. Montel, Sur les suites infinies des fonctions, *Ann. Ecole Norm. sup.* **24** 233–334 (1907).
- [20] C.T. Chuang, *Normal Families of meromorphic Functions*, World Scientific, (1993).
- [21] L. Zalcman, Normal Families: New Perspectives, *Bulletin(New Series) AMS*, **35**, No. 3, 215–230 (1998).